

Knowing for whom the bell tolls: acting locally and thinking globally. Brazil, Latin America and the Global Burden of Diseases, 2015

Sabendo por quem os sinos doam: atuando localmente e pensando globalmente. Brasil, América Latina e a carga global de doenças, 2015

Paulo Andrade Lotufo¹

Faculdade de Medicina da Universidade de São Paulo (FMUSP), São Paulo, SP, Brazil

¹MD, DrPH. Full Professor, Department of Internal Medicine, Faculdade de Medicina da Universidade de São Paulo (FMUSP), São Paulo (SP), Brazil.

“No man is an Iland, intire of it selfe; every man is a peece of the Continent, a part of the maine; if a Clod bee washed away by the Sea, Europe is the lesse, as well as if a Promontorie were, as well as if a Mannor of thy friends or of thine owne were; any mans death diminishes me, because I am involved in Mankinde; And therefore never send to know for whom the bell tolls; It tolls for thee” – John Donne (1572-1623) Devotions upon Emergent Occasions

The Global Burden of Diseases, Injuries and Risk Factors (GBD) study is an initiative from the Institute for Health Metrics and Evaluation of the University of Washington, with unrestricted grants from the Melinda and Bill Gates Foundation. The GBD study has been developing two concepts to enable geographical and temporal trend comparisons. These are the number of years of life lost (YLLs) and number of years of life lived with disability (YLDs). Combination of these two indicators reveals the number of disability-adjusted life-years (DALYs). Moreover, the GBD study is publishing data relating to risk factors. A more detailed description of these indexes can be found in the special issue of *The Lancet* dated October 7, 2016. The data presented below are based on the results of the articles addressing YLL,¹ YLD,² DALY³ and risk factors.⁴

The world can be divided geopolitically in several ways, but there is one grouping that is applied by most agencies: the category of Latin American and Caribbean countries. It represents 50 independent countries and a few colonies with 640 million inhabitants. However, two-thirds of this population and three-quarters of the gross product is concentrated in four countries: Brazil, Mexico, Colombia and Argentina. We will discuss the differences in the GBD indexes among these four countries (**Table 1**).

MORTALITY

The ranking of the five most important causes of YLLs shows how important the burden of coronary heart disease and stroke is, both worldwide and in these four countries, for both sexes. The coronary heart disease (CHD) values were similar among the four countries, but the risk of death due to stroke was significantly higher in Brazil for men and women than in Argentina, Colombia and Mexico.

Two particular causes in Latin American and the Caribbean countries are violence and road traffic, which are ranked within the top five causes of deaths, in contrast to the global data. The road traffic death rate was highest in Brazil (24 per 100,000) followed by Colombia (17), Mexico (17) and Argentina (14). However, the proportion of pedestrian deaths was greater in Colombia (62%) followed by Mexico (44%), Brazil (35%) and Argentina (28%). Violence is a characteristic of Latin American and Caribbean countries: the age-adjusted homicide rates

(per 100,000) in 2015 were 34 in Colombia, 28 in Brazil and 17 in Mexico, but only 6 in Argentina. Another four Latin American and Caribbean countries had higher rates than these: Venezuela, El Salvador, Honduras and Guatemala. The proportion of homicides due to firearms was significantly different: Colombia (78%), Brazil (69%), Mexico (59%) and Argentina (50%).

Another difference observed was the importance of chronic kidney disease (CKD) and diabetes in Mexico. The risk of death due to CKD in Mexico was three times higher than in Argentina, Brazil and Colombia.

YEARS LIVING WITH DISABILITY

Both globally and in these four Latin American and Caribbean countries, an impressive amount of time is lost through disability relating to back and neck disorders, sensory problems, anxiety and depression and skin complaints. However, in contrast to the other countries, diabetes appears as one of the greatest causes of YLDs in Mexico.

DALYS

DALYs are derived from a combination of YLLs and YLDs. They show that CHD is the most important cause in Brazil and Argentina, as also seen worldwide, and that it is the second

biggest cause in Colombia and Mexico. Violence is the top cause in Colombia and diabetes leads the causes of DALYs in Mexico. Neck and back pain and psychiatric disorders have similar impact with regard to DALYs, compared with cardiovascular diseases.

RISK FACTORS

Hypertension is the most important risk factor globally, and in Brazil, Argentina and Colombia, but not in Mexico. The combination of obesity and diabetes is important in all Latin American and Caribbean countries, especially in Mexico, with an association with chronic kidney disease. Smoking was ranked second in the world and in Argentina, fourth in Brazil and fifth in Colombia. Alcohol use was classified as one of the five most important risk factors in Brazil, Colombia and Mexico.

WHAT IS NEW IN GBD 2015?

A relatively large amount of information about mortality and DALYs is available, but the impact of non-lethal conditions on increasing YLDs and DALYs deserves more attention from the Brazilian health authorities. São Paulo Medical Journal dedicated three editorials⁵⁻⁷ and four articles⁸⁻¹¹ to the important topic of lumbar pain. Depression, anxiety and skin disorders need to be more appropriately ranked as priorities for research worldwide,

Table 1. Description of the causes of years of life lost (YLL), years of living with disability (YLD) and disability-adjusted life-years (DALY) for all countries and for Brazil, Argentina, Colombia and Mexico

Global	Brazil	Argentina	Colombia	Mexico
CHD	CHD	CHD	Violence	CHD
Stroke	Violence	LRI	CHD	CKD
LRI	Stroke	Stroke	Road injuries	Diabetes
NN pretem	Road injuries	Road injuries	Congenital	Violence
Diarrhea	NN pretem	Congenital	Stroke	Road injuries
Years of living with disability (YLD)				
Back and neck	Back and neck	Back and neck	Back and neck	Back and neck
Sensory problems	Depression	Depression	Sensory problems	Sensory problems
Depression	Sensory problems	Sensory problems	Depression	Diabetes
Iron	Anxiety	Skin	Skin	Skin
Skin	Skin	Anxiety	Anxiety	Iron
Disability-adjusted life-years (DALY)				
CHD	CHD	CHD	Violence	Diabetes
Stroke	Violence	Back & neck	CHD	CHD
LRI	Stroke	LRI	Back & neck	CKD
Back & neck	Road injuries	Stroke	Sensory problems	Back and neck
NN pretem	Back and neck	Depression	Depression	Sensory problems
Risk factors				
Hypertension	Hypertension	Hypertension	Hypertension	Diabetes
Smoking	Obesity	Smoking	Obesity	Obesity
Diabetes	Diabetes	Obesity	Diabetes	Hypertension
Obesity	Smoking	Diabetes	Alcohol use	Renal dysfunction
Childhood malnutrition	Alcohol use	Dyslipidemia	Smoking	Alcohol use

CHD = coronary heart disease; LRI = lower respiratory infection; CKD = chronic kidney disease; NN pretem = neonatal preterm disorders sensory disabilities.

and the focus of medical care needs to be shifted to primary care, such that it is not managed only by medical specialists.

WHAT ARE THE PRIORITIES FOR ARGENTINA, BRAZIL, COLOMBIA AND MEXICO?

In relation to the epidemiological profile of chronic diseases, GBD 2015 makes it possible to establish five priorities in Latin America and Caribbean countries:

1. **Alcohol intake reduction:** High alcohol consumption is strongly associated with car and motorcycle crashes. Latin America and Caribbean governments are notoriously lenient with regard to punishing individuals who drive under the influence of alcohol beverages. Moreover, deadly interpersonal violence is also related to alcohol abuse.^{12,13}
2. **Hypertension:** Reduction of high blood pressure can reduce racial and social disparities regarding deaths due to cardiovascular diseases. In all countries, it is essential:
 1. to improve awareness, treatment and control of hypertension;
 2. to reduce sodium intake;
 3. to spread the use of automatic sphygmomanometer devices with greater precision and accuracy;^{14,15}
 4. to deliver antihypertensive drugs free of charge (shifting from thiazides to chlorthalidone or indapamide); and
 5. to take special care of individuals with resistant hypertension, who account for 3% of the adult population.¹⁶
3. **CHD secondary prevention:** CHD is the leading cause of death, and the first cause of DALYs in Argentina and Brazil and the second one in Colombia and Mexico. The high prevalence rates of people with CHD implies to amplify national program addressing secondary prevention including smoking quitting and free delivery of aspirin, statins, and angiotensin-converting-enzyme (ACE) inhibitors.¹⁷
4. **Obesity-diabetes prevention:** The prevalence of overweight and obesity in Mexico has been increasing at an alarming rate, with high rates of adverse outcomes relating to diabetes. However, although it is easy to put forward proposals for curbing the obesity epidemic (reduction of calorie intake and increase of physical activity), such proposals are very ineffective. One strategy should be to focus on childhood, so as to avoid obesity among the next generation of teenagers and young adults.¹⁸
5. **CKD screening:** Renal failure in Mexico and Central America leads to high rates of DALYs. This is thought to be due to the high prevalence rate of diabetes in Mexico and of Mesoamerican nephropathy due to heat stress or intoxication with herbicides.^{19,20} Although there is no consensus regarding CKD screening through determination of serum creatinine and urinary albumin, verification of the cost-effectiveness of early diagnosing of CKD is urged.^{21,22}

CONCLUSION

The complexity of the “health-disease” process implies that there is a need to plan and think globally, while acting locally. The most important lesson is that there is only one island: the Earth. And also, the bell tolls for everyone.

REFERENCES

1. GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1459-1544.
2. GBD 2015 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1545-1602.
3. GBD 2015 DALYs and HALE Collaborators. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1603-1658.
4. GBD 2015 Risk Factors Collaborators. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet*. 2016;388(10053):1659-1724.
5. Lotufo PA. From screening-driven medicine to symptom-driven medicine. *Sao Paulo Med J*. 2016;134(5):369-70.
6. Benseñor IJM. Bipedal locomotion, spinal pain and psychiatric disorders. Is this our future? *Sao Paulo Med J*. 2016;134(5):371-2.
7. Suemoto CK. Towards a unified and standardized definition of the frailty phenotype. *Sao Paulo Med J*. 2016;134(5):373-4.
8. Depintor JDP, Bracher ESB, Cabra DMC, Eluf-Neto J. Prevalence of chronic spinal pain and identification of associated factors in a sample of the population of São Paulo, Brazil: cross-sectional study. *Sao Paulo Med J*. 2016;134(5):375-84.
9. Calado LB, Ferriolli E, Moriguti JC, Martinez EZ, Lima NKC. Frailty syndrome in an independent urban population in Brazil (FIBRA study): a cross-sectional populational study. *Sao Paulo Med J*. 2016;134(5):385-92.
10. Tavares DMS, Colamego CG, Pegorari MS, et al. Cardiovascular risk factors associated with frailty syndrome among hospitalized elderly people: a cross-sectional study. *São Paulo Med J*. 2016;134(5):393-99.
11. Orlandini A, Brumini C, Jones A, Natour J. Translation to Brazilian Portuguese, cultural adaptation and reproducibility of the questionnaire “Ankylosing Spondylitis: What do you know?” *Sao Paulo Med J*. 134(5):407-16.
12. Darke S. The toxicology of homicide offenders and victims: A review. *Drug Alcohol Rev*. 2010;29(2):202-15.

13. Kuhns JB, Exum ML, Clodfelter TA, Bottia MC. The prevalence of alcohol-involved homicide offending: a meta-analytic review. *Homicide Studies*. 2013. DOI: 10.1177/1088767913493629. Available from: <http://hsx.sagepub.com/content/early/2013/07/03/1088767913493629.abstract>. Accessed in 2016 (Oct 28).
14. Cloutier L, Daskalopoulou SS, Padwal RS, et al. A New Algorithm for the Diagnosis of Hypertension in Canada. *Can J Cardiol*. 2015;31(5):620-30.
15. Myers MG, Valdivieso M, Kiss A. Use of automated office blood pressure measurement to reduce the white coat response. *J Hypertens*. 2009;27(2):280-6.
16. Lotufo PA, Pereira AC, Vasconcellos PS, et al. Resistant hypertension: risk factors, subclinical atherosclerosis, and comorbidities among adults—the Brazilian Longitudinal Study of Adult Health (ELSA-Brasil). *J Clin Hypertens (Greenwich)*. 2015;17(1):74-80.
17. Lotufo PA. Cardiovascular diseases in Brazil: premature mortality, risk factors and priorities for action. Comments on the preliminary results from the Brazilian National Health Survey (PNS), 2013. *Sao Paulo Med J*. 2015;133(2):69-72.
18. Gortmaker SL, Wang YC, Long MW, et al. Three Interventions That Reduce Childhood Obesity Are Projected To Save More Than They Cost To Implement. *Health Aff (Millwood)*. 2015;34(11):1932-9.
19. Correa-Rotter R, Wesseling C, Johnson RJ. CKD of unknown origin in Central America: the case for a Mesoamerican nephropathy. *Am J Kidney Dis*. 2014;63(3):506-20.
20. Orduñez P, Silva LC. Pesticides and the epidemic of CKD in Central America. *Am J Kidney Dis*. 2014;64(3):477.
21. Moyer VA; US Preventive Services Task Force. Screening for chronic kidney disease: U.S. Preventive Services task force recommendation statement. *Ann Intern Med*. 2012;157(8):567-70.
22. Lotufo PA. Renal disease screening: a potential tool for reducing health inequity. *Sao Paulo Med J*. 2016;134(1):1-2.

Sources of funding: Not declared

Conflict of interest: Not declared

Address for correspondence:

Paulo Andrade Lotufo
Centro de Pesquisa Clínica e Epidemiológica, Hospital Universitário,
Universidade de São Paulo
Av. Prof. Lineu Prestes, 2.565
Butantã — São Paulo (SP) — Brasil
Tel. (+55 11) 3091-9300
E-mail: palotufo@usp.br